

EMMIX-uskew: An R Package for Fitting Mixtures of Multivariate Skew t -distributions via the EM Algorithm

Sharon X. Lee
University of Queensland

Geoffrey J. McLachlan
University of Queensland

Abstract

This paper describes an algorithm for fitting finite mixtures of unrestricted Multivariate Skew t (FM-uMST) distributions. The package **EMMIX-uskew** implements a closed-form expectation-maximization (EM) algorithm for computing the maximum likelihood (ML) estimates of the parameters for the (unrestricted) FM-MST model in R. **EMMIX-uskew** also supports visualization of fitted contours in two and three dimensions, and random sample generation from a specified FM-uMST distribution.

Finite mixtures of skew t -distributions have proven to be useful in modelling heterogeneous data with asymmetric and heavy tail behaviour, for example, datasets from flow cytometry. In recent years, various versions of mixtures with multivariate skew t (MST) distributions have been proposed. However, these models adopted some restricted characterizations of the component MST distributions so that the E-step of the EM algorithm can be evaluated in closed form. This paper focuses on mixtures with unrestricted MST components, and describes an iterative algorithm for the computation of the ML estimates of its model parameters. Its implementation in R is presented with the package **EMMIX-uskew**.

The usefulness of the proposed algorithm is demonstrated in three applications to real data sets. The first example illustrates the use of the main function `fmmst` in the package by fitting a MST distribution to a bivariate unimodal flow cytometric sample. The second example fits a mixture of MST distributions to the Australian Institute of Sport (AIS) data, and demonstrate that **EMMIX-uskew** can provide better clustering results than mixtures with restricted MST components. In the third example, **EMMIX-uskew** is applied to classify cells in a trivariate flow cytometric dataset. Comparisons with other available methods suggests that the **EMMIX-uskew** result achieved a lower misclassification rate with respect to the labels given by benchmark gating analysis.

Keywords: mixture models, skew distributions, multivariate t -distribution, EM algorithm, flow cytometry, R.

1. Introduction

In many practical problems, data are often skewed, heterogeneous, and/or contains outliers. Finite mixture distributions of skewed distributions have become increasingly popular in modelling and analyzing such data. This use of finite mixture distributions to model heterogeneous data has undergone intensive development in the past decades, as witnessed by the numerous applications in various scientific fields such as bioinformatics, cluster analysis, genetics, information processing, medicine, and pattern recognition. For a comprehensive survey on mixture models and their applications see, for example, the monographs by [Everitt and Hand \(1981\)](#), [Titterton et al. \(1985\)](#), [McLachlan and Basford \(1988\)](#), [Lindsay \(1995\)](#), [Böhning \(2000\)](#), [McLachlan and Peel \(2000\)](#), [Frühwirth-Schnatter \(2006\)](#), and [Mengersen et al. \(2011\)](#), and also the papers by [Banfield and Raftery \(1993\)](#) and [Fraley and Raftery \(1999\)](#).

In recent years, finite mixtures of skew t -distributions have been exploited as an effective tool in modelling high-dimensional multimodal and asymmetric dataset; see, for example, [Pyne et al. \(2009a\)](#) and [Frühwirth-Schnatter and Pyne \(2010\)](#). Following the introduction of the skew-normal (SN) distribution by [Azzalini \(1985\)](#), several authors have studied skewed extensions of the t -distribution. Finite mixture models with multivariate skew t (MST) components was first proposed by [Pyne et al. \(2009a\)](#) in a study of an automated approach to the analysis of flow cytometry data. [Wang \(2009\)](#) has given a package **EMMIX-skew** for the implementation in R ([R Development Team 2011](#)) of their algorithm. More recently, [Basso et al. \(2010\)](#) studied a class of mixture models where the components densities are scale mixtures of univariate skew-normal distributions, known as the skew-normal/independent (SNI) family of distributions, which include the (univariate) skew-normal and skew t -distributions as special cases. This work was later extended to the multivariate case in [Cabral et al. \(2012\)](#), and was implemented in an R package **mixsmsn**. However, in these characterizations, restrictions were imposed on the component skew t -distributions in order to obtain manageable analytical expressions for the conditional expectations involved in the E-step of the EM algorithm. These versions of the skew t -distributions are known as the ‘restricted’ form of the MST distribution; see [Lee and McLachlan \(2012a\)](#) for further discussions on this.

In this paper, we present a fitting algorithm for the unrestricted skew t -mixture model. We show that an EM algorithm can be implemented exactly without restricting the characterizations of the component MST distributions. Closed form expressions can be obtained for the E-step conditional expectations by recognizing that they can be formulated as moments of a multivariate non-central truncated t -variate, which can be further expressed in terms of central t -distributions. The algorithm is implemented in R in the package **EMMIX-uskew**.

The package **EMMIX-uskew** consists of three main functions: `fmmst`, `rfmmst`, and `contour.fmmst`. The main routine `fmmst` fits a mixture of unrestricted MST (uMST) distributions using an EM algorithm described in Section 3. The function `rfmmst` generates random samples from mixtures of uMST distributions. For a user friendly visualisation of the fitted models, `fmmst.contour` provides 2D contour maps of the fitted bivariate densities and 3D displays with interactive viewpoint navigation facility for trivariate densities.

The remainder of this paper is organized as follows. Section 2 provides a brief description of the uMST distribution and defines the FM-uMST model. Section 3 presents an EM algorithm for fitting the FM-uMST model. In the next section, an explanation of how to fit, visualize, and interpret the FM-uMST models using **EMMIX-uskew** is presented. For illustration purposes, **EMMIX-uskew** is applied to three applications and comparisons are made with some restricted

FM-MST models and other clustering methods. Finally, we conclude with a brief summary of our results.

2. Finite mixtures of multivariate skew t -distributions

We begin by defining the (unrestricted) multivariate skew t -density. Let $\mathbf{Y}$ be a p -dimensional random vector. Then $\mathbf{Y}$ is said to follow a p -dimensional unrestricted skew t -distribution (Sahu *et al.* 2003) with $p \times 1$ location vector $\boldsymbol{\mu}$, $p \times p$ scale matrix $\boldsymbol{\Sigma}$, $p \times 1$ skewness vector $\boldsymbol{\delta}$, and (scalar) degrees of freedom ν , if its probability density function (pdf) is given by

$$f_p(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\delta}, \nu) = 2^p t_{p,\nu}(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Omega}) T_{p,\nu+p}(\mathbf{y}^*; \mathbf{0}, \boldsymbol{\Lambda}), \quad (1)$$

where

$$\begin{aligned} \boldsymbol{\Delta} &= \text{diag}(\boldsymbol{\delta}), \\ \boldsymbol{\Omega} &= \boldsymbol{\Sigma} + \boldsymbol{\Delta}^2, \\ \mathbf{y}^* &= \mathbf{q} \sqrt{\frac{\nu + p}{\nu + d(\mathbf{y})}}, \\ \mathbf{q} &= \boldsymbol{\Delta} \boldsymbol{\Omega}^{-1}(\mathbf{y} - \boldsymbol{\mu}), \\ d(\mathbf{y}) &= (\mathbf{y} - \boldsymbol{\mu})^T \boldsymbol{\Omega}^{-1}(\mathbf{y} - \boldsymbol{\mu}), \\ \boldsymbol{\Lambda} &= \mathbf{I}_p - \boldsymbol{\Delta} \boldsymbol{\Omega}^{-1} \boldsymbol{\Delta}. \end{aligned}$$

Here the operator $\text{diag}(\boldsymbol{\delta})$ denotes a diagonal matrix with diagonal elements specified by the vector $\boldsymbol{\delta}$. Also, we let $t_{p,\nu}(\cdot; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ be the p -dimensional t -distribution with location vector $\boldsymbol{\mu}$, scale matrix $\boldsymbol{\Sigma}$, and degrees of freedom ν , and $T_{p,\nu}(\cdot; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ the corresponding (cumulative) distribution function. The notation $\mathbf{Y} \sim \text{uMST}_{p,\nu}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\delta})$ will be used. Note that when $\boldsymbol{\delta} = \mathbf{0}$, (1) reduces to the symmetric t -density $t_{p,\nu}(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma})$. Also, when $\nu \rightarrow \infty$, we obtain the (unrestricted) skew normal distribution.

Various versions of the multivariate skew t -density have been proposed in recent years. It is worth noting that the versions considered by Azzalini and Dalla Valle (1996), Gupta (2003), and Lachos *et al.* (2010), among others, are different from (1). These versions are simpler in that the skew t -density is defined in terms involving only the univariate t -distribution function instead of the multivariate form of the latter as used in (1). These simplified characterizations have the advantage of having closed form expressions for the conditional expectations that have to be calculated on the E-step. The reader is referred to Lee and McLachlan (2012a,b) for a discussion on different forms of skew t -distributions. We shall adopt the unrestricted form (1) of the MST distribution here as proposed by Sahu *et al.* (2003), and describe a computationally efficient EM algorithm for fitting such model.

A g -component finite mixture of uMST distributions has density given by

$$f(\mathbf{y}; \boldsymbol{\Psi}) = \sum_{h=1}^g \pi_h f_p(\mathbf{y}; \boldsymbol{\mu}_h, \boldsymbol{\Sigma}_h, \boldsymbol{\delta}_h, \nu_h), \quad (2)$$

where $f_p(\mathbf{y}; \boldsymbol{\mu}_h, \boldsymbol{\Sigma}_h, \boldsymbol{\delta}_h, \nu_h)$ denotes the h th uMST component of the mixture model as defined by (1), with location parameter $\boldsymbol{\mu}_h$, scale matrix $\boldsymbol{\Sigma}_h$, skew parameter $\boldsymbol{\delta}_h$, and degrees

of freedom ν_h . The mixing proportions π_h satisfy $\pi_h \geq 0$ ($h = 1, \dots, g$) and $\sum_{h=1}^g \pi_h = 1$. We shall denote the model defined by (2) by the FM-uMST (finite mixture of uMST) distributions. Let Ψ contain all the unknown parameters of the FM-uMST model; that is, $\Psi = (\pi_1, \dots, \pi_{g-1}, \theta_1^T, \dots, \theta_g^T)^T$ where now θ_h consists of the unknown parameters of the h th component density function. The density values for a uMST and FM-uMST distribution can be evaluated using the functions `dmst` and `dfmmst` in **EMMIX-uskew**.

Random samples of uMST variates can be generated by adopting a stochastic representation of (1) (Lin 2010). If $\mathbf{Y} \sim \text{uMST}_{p,\nu}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\delta})$, then

$$\mathbf{Y} = \boldsymbol{\mu} + \frac{1}{\sqrt{w}} \boldsymbol{\Delta} |\mathbf{U}_1| + \frac{1}{\sqrt{w}} \mathbf{U}_0, \quad (3)$$

where the random variables

$$\mathbf{U}_0 \sim N_p(\mathbf{0}, \boldsymbol{\Sigma}), \quad (4)$$

$$\mathbf{U}_1 \sim N_p(\mathbf{0}, \mathbf{I}_p), \quad (5)$$

$$w \sim \text{gamma}\left(\frac{\nu}{2}, \frac{\nu}{2}\right), \quad (6)$$

are independent, and $\text{gamma}(\alpha, \beta)$ denotes the gamma distribution with shape and scale parameters given by α and β respectively. Sampling of uMST and FM-uMST variates are implemented in **EMMIX-uskew** in the `rmst` and `rfmmst` functions, respectively.

3. The EMMIX-uskew algorithm

From (3) to (6), the uMST distribution admits a convenient hierarchical characterization that facilitates the computation of the maximum likelihood estimator (MLE) of the unknown model parameters using the EM algorithm, namely,

$$\begin{aligned} \mathbf{Y}_j | \mathbf{u}_j, w_j &\sim N_p\left(\boldsymbol{\mu} + \boldsymbol{\Delta} \mathbf{u}_j, \frac{1}{w_j} \boldsymbol{\Sigma}\right), \\ \mathbf{U}_j | w_j &\sim HN_p\left(\mathbf{0}, \frac{1}{w_j} \mathbf{I}_p\right), \\ W_j &\sim \text{gamma}\left(\frac{\nu}{2}, \frac{\nu}{2}\right), \end{aligned}$$

where $\boldsymbol{\Delta}_h = \text{diag}(\boldsymbol{\delta}_h)$, $HN_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes the p -dimensional half-normal distribution with location parameter $\boldsymbol{\mu}$ and scale matrix $\boldsymbol{\Sigma}$, and $\boldsymbol{\pi} = (\pi_1, \dots, \pi_g)^T$.

3.1. Fitting of FM-uMST model via the EM algorithm

To formulate the estimation of the unknown parameters as an incomplete-data problem in the EM framework, we introduce a set of latent component labels $\mathbf{z}_j = (z_{1j}, \dots, z_{gj})$ ($j = 1, \dots, n$) in addition to the unobservable variables $\mathbf{u}_j$ and w_j , where each element z_{hj} is a zero-one indicator variable with $z_{hj} = 1$ if $\mathbf{y}_j$ belongs to the h th component, and zero otherwise. Thus, $\sum_{h=1}^g z_{hj} = 1$ ($j = 1, \dots, n$). It follows that the random vector $\mathbf{Z}_j$ corresponding to $\mathbf{z}_j$ follows a multinomial distribution with one trial and cell probabilities $\pi_1, \dots, \pi_g$; that is, $\mathbf{Z}_j \sim \text{Mult}_g(1; \pi_1, \dots, \pi_g)$.

The complete-data log likelihood function can be factored into the marginal densities of the $\mathbf{z}_j$, the conditional densities of the w_j given $\mathbf{z}_j$, and the conditional densities of the $\mathbf{y}_j$ given $\mathbf{u}_j$, w_j , and $\mathbf{z}_j$. Accordingly, the complete-data log likelihood is given by

$$\log L_c(\Psi) = \log L_{1c}(\Psi) + \log L_{2c}(\Psi) + \log L_{3c}(\Psi), \quad (7)$$

where

$$\begin{aligned} L_{1c}(\Psi) &= \sum_{h=1}^g \sum_{j=1}^n z_{hj} \log(\pi_h), \\ L_{2c}(\Psi) &= \sum_{h=1}^g \sum_{j=1}^n z_{hj} \left[\left(\frac{\nu_h}{2} \right) \log \left(\frac{\nu_h}{2} \right) + \left(\frac{\nu_h}{2} + p - 1 \right) \log(w_j) \right. \\ &\quad \left. - \log \Gamma \left(\frac{\nu_h}{2} \right) - \left(\frac{w_j}{2} \right) \nu_h \right], \\ L_{3c}(\Psi) &= \sum_{h=1}^g \sum_{j=1}^n z_{hj} \left\{ -p \log(2\pi) - \frac{1}{2} \log |\Sigma_h| \right. \\ &\quad \left. - \frac{w_j}{2} \left[d_h(\mathbf{y}_j) + (\mathbf{u}_j - \mathbf{q}_{hj})^T \Lambda_h^{-1} (\mathbf{u}_j - \mathbf{q}_{hj}) \right] \right\}, \end{aligned} \quad (8)$$

and where

$$\begin{aligned} d_h(\mathbf{y}_j) &= (\mathbf{y}_j - \boldsymbol{\mu}_h)^T \boldsymbol{\Omega}_h^{-1} (\mathbf{y}_j - \boldsymbol{\mu}_h), \\ \mathbf{q}_{hj} &= \boldsymbol{\Delta}_h \boldsymbol{\Omega}_h^{-1} (\mathbf{y}_j - \boldsymbol{\mu}_h), \\ \Lambda_h &= \mathbf{I}_p - \boldsymbol{\Delta}_h \boldsymbol{\Omega}_h^{-1} \boldsymbol{\Delta}_h, \\ \boldsymbol{\Omega}_h &= \Sigma_h + \boldsymbol{\Delta}_h^2. \end{aligned}$$

Here Ψ contains all the unknown parameters of the FM-uMST model.

The implementation of the EM algorithm requires alternating repeatedly the E- and M-steps until convergence in the case where the changes in the log likelihood values are less than some specified small value. The E-step calculates the expectation of the complete-data log likelihood given the observed data $\mathbf{y}$ using the current estimate of the parameters, known as the Q -function, given by

$$Q(\Psi; \Psi^{(k)}) = E_{\Psi^{(k)}} \{ \log L_c(\Psi) \mid \mathbf{y} \}.$$

The M-step then maximizes the Q -function with respect to the parameters Ψ .

On the $(k+1)$ th iteration, the E-step requires the calculation of the conditional expectations

$$e_{1,j}^{(k)} = E_{\boldsymbol{\theta}^{(k)}} (W_j \mid \mathbf{y}_j), \quad (9)$$

$$\mathbf{e}_{2,j}^{(k)} = E_{\boldsymbol{\theta}^{(k)}} (W_j \mathbf{U}_j \mid \mathbf{y}_j), \quad (10)$$

$$\mathbf{e}_{3,j}^{(k)} = E_{\boldsymbol{\theta}^{(k)}} (W_j \mathbf{U}_j \mathbf{U}_j^T \mid \mathbf{y}_j). \quad (11)$$

The posterior probability of membership of the h th component by $\mathbf{y}_j$, using the current estimate $\Psi^{(k)}$ for Ψ , is given using Bayes' Theorem by

$$\tau_{hj}^{(k)} = \frac{\pi_h^{(k)} f_p(\mathbf{y}_j; \boldsymbol{\mu}_h^{(k)}, \Sigma_h^{(k)}, \boldsymbol{\delta}_h^{(k)}, \nu_h^{(k)})}{\sum_{i=1}^g \pi_i^{(k)} f_p(\mathbf{y}_j; \boldsymbol{\mu}_i^{(k)}, \Sigma_i^{(k)}, \boldsymbol{\delta}_i^{(k)}, \nu_i^{(k)})}. \quad (12)$$

It can be shown that the conditional expectations $e_{1,j}^{(k)}$, $e_{2,j}^{(k)}$, and $e_{3,j}^{(k)}$ are given by

$$e_{1,hj}^{(k)} = \left(\frac{\nu_h^{(k)} + p}{\nu_h^{(k)} + d_h^{(k)}(\mathbf{y}_j)} \right) \frac{T_{p, \nu_h^{(k)} + p + 2} \left(\mathbf{q}_{hj}^{(k)} \sqrt{\frac{\nu_h^{(k)} + p + 2}{\nu_h^{(k)} + d_h^{(k)}(\mathbf{y}_j)}}; \mathbf{0}, \mathbf{\Lambda}_h^{(k)} \right)}{T_{p, \nu_h^{(k)} + p} \left(\mathbf{y}_{hj}^{*(k)}; \mathbf{0}, \mathbf{\Lambda}_h^{(k)} \right)}, \quad (13)$$

$$e_{2,hj}^{(k)} = e_{1,hj}^{(k)} E(\mathbf{X}), \quad (14)$$

and

$$e_{3,hj}^{(k)} = e_{1,hj}^{(k)} E(\mathbf{X} \mathbf{X}^T), \quad (15)$$

where $\mathbf{X}$ is a p -dimensional t -variate truncated to the positive hyperplane $\mathbb{R}^+$, which is distributed as

$$\mathbf{X} \sim tt_{p, \nu_h^{(k)} + p + 2} \left(\mathbf{q}_{hj}^{(k)}, \left(\frac{\nu_h^{(k)} + d_h^{(k)}(\mathbf{y}_j)}{\nu_h^{(k)} + p + 2} \right) \mathbf{\Lambda}_h^{(k)}; \mathbb{R}^+ \right). \quad (16)$$

The truncated moments $E(\mathbf{X})$ and $E(\mathbf{X} \mathbf{X}^T)$ can be swiftly evaluated by noting that they can be expressed in terms of the distribution function of a (non-truncated) multivariate central t -random vector; Lee and McLachlan (2011, 2012a). Recently, Ho *et al.* (2012) have considered the moments of the doubly truncated multivariate t -distribution, but their result corresponding to (16) would appear to be incorrect (see Lee and McLachlan (2012a)).

The $(k+1)$ th M-step consists of maximization of the Q -function with respect to Ψ . It follows that an updated estimate of the unknown parameters of the FM-uMST model is given by

$$\mu_h^{(k)} = \frac{\sum_{j=1}^n \tau_{hj}^{(k)} [e_{1,hj}^{(k)} \mathbf{y}_j - \Delta_h^{(k)} e_{2,hj}^{(k)}]}{\sum_{j=1}^n \tau_{hj}^{(k)} e_{1,hj}^{(k)}}, \quad (17)$$

$$\delta_h^{(k+1)} = \left(\Sigma_h^{(k)-1} \odot \sum_{j=1}^n \tau_{hj}^{(k)} e_{3,hj}^{(k)} \right)^{-1} \text{diag} \left(\Sigma_h^{(k)-1} \sum_{j=1}^n \tau_{hj}^{(k)} (\mathbf{y}_j - \mu_h^{(k+1)}) e_{2,hj}^{(k)T} \right), \quad (18)$$

and

$$\begin{aligned} \Sigma_h^{(k+1)} = & \frac{1}{\sum_{j=1}^n \tau_{hj}^{(k)}} \sum_{j=1}^n \tau_{hj}^{(k)} \left[\Delta_h^{(k+1)} e_{3,hj}^{(k)T} \Delta_h^{(k+1)T} - (\mathbf{y}_j - \mu_h^{(k+1)}) e_{2,hj}^{(k)T} \Delta_h^{(k+1)} \right. \\ & \left. - \Delta_h^{(k+1)} e_{2,hj}^{(k)} (\mathbf{y}_j - \mu_h^{(k+1)})^T + (\mathbf{y}_j - \mu_h^{(k+1)}) (\mathbf{y}_j - \mu_h^{(k+1)})^T e_{1,hj}^{(k)} \right], \end{aligned} \quad (19)$$

where $\odot$ denotes element-wise matrix product. Note that (18) and also (16) are given incorrectly in Lee and McLachlan (2011).

An update $\nu_h^{(k+1)}$ of the degrees of freedom is obtained by solving iteratively the equation

$$\log \left(\frac{\nu_h^{(k+1)}}{2} \right) - \psi \left(\frac{\nu_h^{(k+1)}}{2} \right) = \frac{\sum_{j=1}^n \tau_{hj}^{(k)} \left[\log \left(\frac{\nu_h^{(k)} + d_h^{(k)}(\mathbf{y}_j)}{2} \right) - \psi \left(\frac{\nu_h^{(k)} + p}{2} \right) + \frac{\nu_h^{(k)} + p}{\nu_h^{(k)} + d_h^{(k)}(\mathbf{y}_j)} \right]}{\sum_{j=1}^n \tau_{hj}^{(k)}},$$

where $\psi(\cdot)$ is the Digamma function.

This algorithm is implemented as the `fmmst` function in **EMMIX-uskew**.

3.2. Choosing initial values

It is important to obtain suitable initial values in order for `fmmst` to converge quickly. In **EMMIX-uskew**, starting values for the model parameters are based on an initial clustering given by k -means. Twenty attempts of k -means are performed, and the starting component labels $\mathbf{z}_j^{(0)}$ ($j = 1, \dots, n$) are initialized according to the clustering result with the highest relative log likelihood (see Lee and McLachlan (2012a)). The other parameters are initialized as follows:

$$\begin{aligned}\Sigma^{(0)} &= \mathbf{S}_h - (a - 1) \text{diag}(\mathbf{s}_h), \\ \boldsymbol{\delta}^{(0)} &= \text{sign}(\boldsymbol{\gamma}_h) \sqrt{\frac{(1-a)\pi}{\pi-2}} \mathbf{s}_h^*, \\ \boldsymbol{\mu}^{(0)} &= \bar{\mathbf{y}} - \sqrt{\frac{2}{\pi}} \boldsymbol{\delta}^{(0)}, \\ \nu^{(0)} &= 40,\end{aligned}$$

where $\mathbf{S}_h$ is the sample covariance of the h th component, and where $\boldsymbol{\gamma}_h$ is the sample skewness of the h th component, whose i th element is given by

$$\gamma_i = \frac{n^{-1} \sum_{j=1}^n (y_{ij} - \mu_i)^3}{\left(n^{-1} \sum_{j=1}^n (y_{ij} - \mu_i)^2\right)^{\frac{3}{2}}} \quad (i = 1, \dots, p),$$

and where y_{ij} denotes the i th element of the j th observation, and μ_i is the i th element of $\boldsymbol{\mu}$. Here, $\mathbf{s}_h$ denotes the vector created by extracting the main diagonal of $\mathbf{S}_h$, and the vector $\mathbf{s}_h^*$ is created by taking the square root of each element in $\mathbf{s}_h$. The scalar a is varied systematically across the interval $(0, 1)$ to search for a (relatively) optimal set of starting values for the model parameters.

3.3. Stopping rule

EMMIX-uskew adopts the traditional stopping criteria which is based on the relative change in the size of the log likelihood. The algorithm is terminated when the relative difference in log likelihood values between two successive iteration is less than a specified tolerance ϵ , that is, the EM algorithm is stopped if

$$\frac{|L^{(k+1)} - L^{(k)}|}{|L^{(k+1)}|} < \epsilon,$$

where $L^{(k+1)}$ denotes the log likelihood value at the $(k+1)$ th iteration. The default tolerance is $\epsilon = 10^{-6}$, but the user can specify a different value.

4. Using the EMMIX-uskew package

The parameters of the FM-uMST model in **EMMIX-uskew** are specified as a list structure containing the elements described in Table 1. The parameters $\boldsymbol{\mu}$, $\boldsymbol{\Sigma}$, and $\boldsymbol{\delta}$ are each implemented as a list of $\mathbf{g}$ matrices, where $\mathbf{g}$ is the number of components in the fitted model. For example, `mu[[2]]` is a $p \times 1$ matrix representing $\boldsymbol{\mu}_2$. Each `sigma[[h]]` ($h = 1, \dots, g$) is a $p \times p$ matrix representing the symmetric positive definite scale matrix of the h th component. The parameters `nu` and `pro` are g by 1 arrays, representing the vector of degrees of freedom and the vector of mixing proportions for each component, respectively.

parameter	R syntax	Dimensions	Description
$\boldsymbol{\mu}$	<code>mu</code>	$p \times 1 \times g$	the location parameter
$\boldsymbol{\Sigma}$	<code>sigma</code>	$p \times p \times g$	the scale matrix
$\boldsymbol{\delta}$	<code>delta</code>	$p \times 1 \times g$	the skewness parameter
ν	<code>dof</code>	$g \times 1$	the degrees of freedom
π	<code>pi</code>	$g \times 1$	the mixing proportions

Table 1: Structure of the model parameters in **EMMIX-uskew**.

The probability density function of a multivariate skew t -distribution is calculated by the `dmst` function. The parameter `dat` is an $n \times p$ matrix, containing the coordinates of the n point(s) at which the density is to be evaluated. The following command will return a vector of n density values.

```
dmst(dat, mu, sigma, delta, dof)
```

For a FM-uMST density, the function `dfmmst` can be used.

```
dfmmst(dat, mu, sigma, delta, dof, pro)
```

4.1. Generating samples from a FM-uMST distribution

Consider generating a random sample of n p -dimensional uMST observations, with location parameter $\boldsymbol{\mu}$, scale matrix $\boldsymbol{\Sigma}$, skewness parameter $\boldsymbol{\delta}$, and degrees of freedom ν . The function `rfmmst` supports two types of inputs – the parameters can be passed as separate arguments, or as a single list argument `known` with elements as specified in Table 1:

```
rfmmst(g, n, mu, sigma, delta, dof, pro, known=NULL, ...)
```

As an example, suppose that $\boldsymbol{\mu} = (1, 2)^T$, $\boldsymbol{\Sigma}$ is the identity matrix, $\boldsymbol{\delta} = (-1, 1)^T$, and $\nu = 4$. Then the following command will generate a random sample of 500 observations from the $uMST_{2,4}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\delta})$ distribution,

```
R> rfmmst(1, 500, c(1, 2), diag(2), c(-1, 1), 4, 1)
```

To generate a mixture of uMST random samples, the above command can be issued. Alternatively, the parameters can be specified in a list structure (Table 1) `obj` as follows:

```

R> obj <- list()
R> obj$mu <- list(c(17,19), c(5,22), c(6,10))
R> obj$sigma <- list(diag(2), matrix(c(2,0,0,1),2), matrix(c(3,7,7,24),2))
R> obj$delta <- list(c(3,1.5), c(5,10), c(2,0))
R> obj$dof <- c(1, 2, 3)
R> obj$pro <- c(0.25, 0.25, 0.5)
R> rfmmst(3, 500, known=obj)

```

An output of the `rfmmst` function consists of $p + 1$ columns. The first p columns are the coordinates of the generated sample. The last column indicates from which component each data point is generated. Executing the above command will generate an output similar to the following:

```

      [,1]      [,2]      [,3]
[1,] 19.91520 20.48515      1
[2,] 72.81161 33.41381      1
[3,] 17.02193 23.29119      1
[4,] 23.53926 19.27946      1
[5,] 16.85195 21.21340      1
[6,] 18.01906 18.16612      1
[7,] 22.23609 21.12174      1
[8,] 44.65444 28.23259      1
[9,] 18.18883 26.72330      1
[10,] 20.18908 18.97005      1
... rest omitted ...

```

4.2. Fitting a single multivariate skew t -distribution

To fit a specified FM-uMST model, the core function in **EMMIX-uskew**, `fmmst`, is used. This implements the algorithm described in Section 3. A typical function call of `fmmst` is:

```
fmmst(g, dat, initial=NULL, known=NULL, itmax=100, eps=1e-6, debug=TRUE)
```

The main arguments used within this function are:

- **g**: a scalar that specifies the number of uMST components to be fitted.
- **dat**: an $n \times p$ matrix containing the data.
- **initial**: a list that specifies the initial values used to start the algorithm.
- **known**: a list that specifies any model parameters that are known and so not required to be estimated.
- **itmax**: a scalar that specifies the maximum number of iterations to be used.
- **eps**: a scalar that specifies the termination criterion of the EM algorithm loop.

Note that if the initial values of the model parameters are provided by the user, the argument `initial` is expected to be structured as described in Table 1. Similarly, `known` is expected to have the same structure. When `initial=NULL`, `fmmst` will generate a set of initial values using the procedure described in Section 3.2. Any parameters specified in `known` are taken as known parameters and hence are not estimated by `fmmst`. There is no need to specify the values of all the parameters in `initial` and `known` when only some of the parameters are known. Parameters that are not specified during the function call are estimated by `fmmst`. The termination criteria for the **EMMIX-uskew** algorithm is controlled by the parameters `itmax` and `eps`. The EM loop terminates when either one of the two criterion is satisfied, whichever occurs first: (a) the EM loop reaches `itmax` iterations (default is 100 iterations), or (b) the relative difference between the current log likelihood value and that from the previous iteration is smaller than `eps` (default is $1e-6$). The last argument of `fmmst` is `debug`. When the option `debug` is set to `TRUE` (default), `fmmst` prints the log likelihood value at each iteration and displays a summary of the parameters of the fitted model after termination. To turn off debug mode, simply set `debug=FALSE`. For further details of the arguments of `fmmst`, the reader is referred to the documentation of `fmmst`. This can be accessed by typing `?fmmst` at the R command prompt.

We consider now the T-cell phosphorylation dataset (Maier *et al.* 2007) as an example of asymmetrically distributed data, available from Pyne *et al.* (2009b). The data contain measurements of blood samples stained with four antibodies, CD4, CD45RA, SLP76, and ZAP70. For illustration, we randomly select 500 observations and focus on two of the variables, CD4 and ZAP70. To fit a MST model to this bivariate Lymphoma dataset, under the default settings, the following command is issued:

```
R> set.seed(12345)
R > data("Lympho")
R > LymphoSample <- Lympho[sample(1:nrow(Lympho), 500),]
R > Fit <- fmmst(1, LymphoSample)
```

A summary of the output of the fitted model can be obtained using the `summary` function. This prints the values of the fitted model parameters for each component. For a fitted uMST model, the weighting proportion (which is 1) is not printed. The following output shows a typical summary of a fitted single component uMST model.

```
R > summary(Fit)
Finite Mixture of Multivariate Skew t-Distribution
with 1 component

Mean:
      [,1]
[1,] 4.808245
[2,] 5.500559

Scale matrix:
[[1]]
      [,1]      [,2]
```

```
[1,] 0.06778378 0.03721489
[2,] 0.03721489 0.04811898
```

Skewness parameter:

```
      [,1]
[1,] -0.7082174
[2,] -0.7990700
```

Degrees of freedom:

```
5.851434
```

```
> summary(Fit)
```

```
Finite Mixture of Multivariate Skew t-Distributions
with 1 component
```

Mean:

```
      [,1]
[1,] 4.808245
[2,] 5.500559
```

Scale matrix:

```
[[1]]
      [,1]      [,2]
[1,] 0.06778378 0.03721489
[2,] 0.03721489 0.04811898
```

Skewness parameter:

```
      [,1]
[1,] -0.7082174
[2,] -0.7990700
```

Degrees of freedom:

```
5.851434
```

To view a more detailed output of the `fmmst` function, the `print` function is called. This outputs a list containing 11 elements. The first five elements give the estimates of the parameters of the fitted FM-uMST model, as described in Table 1.

The posterior probability of component membership is given by the output argument `tau`, a $g \times n$ matrix where the rows corresponds to the component number. The final partition of each data point, based on `tau`, is stored as `clusters`. The value of the log likelihood function, evaluated with the current parameter estimates, is given by `loglik`. The last two arguments `aic` and `bic` are the values of the Akaike information criterion (AIC) and the Bayes information criterion (BIC), respectively. The following output shows an excerpt from the second part of the `print` output of the fitted model.

```

R> print(Fit)
Finite Mixture of Multivariate Skew t-Distributions
with 1 component
... first five components omitted ...

$tau
      [,1] [,2] [,3] [,4] [,5] [,6] [,7] [,8] [,9] [,10] [,11] [,12] [,13] [,14]
[1,]      1      1      1      1      1      1      1      1      1      1      1      1      1      1
... rest omitted ...

$clusters
[1]      1      1      1      1      1      1      1      1      1      1      1      1      1      1
... rest omitted ...

$ik
[1] -880.7115

$aic
[1] 1777.423

$bic
[1] 1811.14

```

4.3. Fitting mixtures of multivariate skew t -distributions

This section presents an illustration of fitting a mixture of unrestricted skew t -distributions to some bivariate bimodal asymmetric data. We consider the Australian Institute of Sport (AIS) data from [Cook and Weisberg \(1994\)](#), where thirteen body measurements on 102 male and 100 female athletes were recorded. In this demonstration, we consider the clustering of the data with a two component skew t -mixture model based on the two variables Height and Body fat. By setting `debug=TRUE`, we can examine the value of the log likelihood function at each iteration.

```
R> Fit2 <- fmmst(2, ais[,c(2,12)], debug=TRUE)
```

```

Finite Mixture of Multivariate Skew t-Distributions
with 2 components

```

```

-----
Iteration 1 : loglik = -1372.711
Iteration 2 : loglik = -1370.495
Iteration 3 : loglik = -1368.773
Iteration 4 : loglik = -1367.392
Iteration 5 : loglik = -1366.251
... rest omitted ...
-----

```

```
Iteration 100 : loglik = -1343.541
```

```
Component means:
```

```
      [,1]      [,2]
[1,] 181.91720 181.448906
[2,] 13.67975  5.814277
```

```
Component scale matrices:
```

```
[[1]]
      [,1]      [,2]
[1,] 26.32964 18.75208
[2,] 18.75208 16.12990
```

```
[[2]]
      [,1]      [,2]
[1,] 61.598537 2.3177048
[2,]  2.317705 0.1515253
```

```
Component skewness parameters:
```

```
      [,1]      [,2]
[1,] -9.582015 3.591328
[2,]  5.975328 5.729339
```

```
Component degrees of freedom:
```

```
60.03386 28.94895
```

```
Component mixing proportions:
```

```
0.4102178 0.5897822
```

We compare the results with two other model-based clustering methods provided by the package **mixsmsn** (Prates *et al.* 2011) and **EMMIX-skew** (Wang 2009). As mentioned previously, these two models are based on mixture of restricted versions of the multivariate skew t -distributions. The first model adopts the skew-normal/independent skew t -distribution (Cabral *et al.* 2012) as its component densities, which is equivalent to the restricted skew t -distribution (Pyne *et al.* 2009a) used in the second model. However, it should be noted that, in the ECME algorithm implemented in the package **mixsmsn**, the component degrees of freedom are constrained to be the same. A comparison of the table of cluster labels to the true class labels (given by `ais$Sex` in this example) reveals that the FM-uMST model has a higher number of correct allocations (183 compared to 161 and 157 given by **mixsmsn** and **EMMIX-skew** respectively). Thus, the unrestricted FM-uMST model in **EMMIX-uskew** gives a more accurate clustering in this case.

```
R> library("mixsmsn")
R> Fit3 <- smsn.mmix(ais[c(2,12)], g=2, family="Skew.t", group=TRUE)
R> Fit4 <- EmSkew(ais[c(2,12)], 2, "mst", debug=FALSE)
R> table(ais$Sex, Fit3$group)
  1  2
```

```

0 91 11
1 29 71
R> table(ais$Sex, Fit4$clust)
  1  2
0 89 13
1 32 68
R> table(ais$Sex, Fit2$clusters)
  1  2
0  5 97
1 86 14

```

4.4. Discriminant analysis

Discriminant analysis based on a specified FM-uMST model can be performed using the `fmmstDA` function.

```
fmmstDA(g, dat, model)
```

The data in `dat` is assigned to the cluster corresponding to the component of the FM-uMST model with the highest posterior probability. Specifications of the model parameters must be provided in `model`, which is typically an output from `fmmst`. The following commands shows an example using `fmmstDA`. A random sample of FM-uMST variables is generated from `rfmmst`, the first part of which is used as training set, and the second is a testing set. The FM-uMST model fitted to the training set is then used for classifying the data in the testing set.

```

R> set.seed(732)
R> X <- rfmmst(3, 200, known=obj)
R> Ind <- sample(1:nrow(X), 175)
R> train <- X[Ind,]
R> test <- X[-Ind,]
R> trainmodel <- fmmst(3, train[,1:2])
R> fmmstDA(3, test[,1:2], trainmodel)
R> results <- fmmstDA(3, test[,1:2], trainmodel)
R> table(test[,3], results)
  results
    1  2  3
1  0  6  0
2  0  0  5
3 13  0  1

```

4.5. Visualization of fitted contours

The **EMMIX-uskew** package supports visualization of the contours of a FM-uMST model in 2D and 3D. The plots are generated by the functions `fmmst.contour.2d` and `fmmst.contour.3d`,

```

fmmst.contour.2d(dat, model, grid=50, drawpoints=TRUE, clusters=NULL,
  levels=10, component=NULL, map=c("scatter", "heat", "cluster"), ...)
fmmst.contour.3d(dat, model, grid=20, drawpoints=TRUE, clusters=NULL,
  levels=0.9, component=NULL, ...)

```

In `fmmst.contour.2d` (`fmmst.contour.3d`), the first argument `dat` is a matrix of coordinates with two (three) columns. The second argument `model` is a list of at least six elements: the five model parameters, and a vector of cluster labels `clusters`. The grid size is determined by `grid`. By default, the data points are included in the plot. If only the contour is required, the option `drawpoints=FALSE` should be set. When including the points in a plot, `clusters` specifies the component labels of each point according to which the data points will be coloured. The argument `levels` is either an integer specifying the number of contour lines to be plotted, or a vector of quantile values. For `fmmst.contour.3d`, only the 90th percentile contour is plotted by default. If more contours are required, the argument `levels` should be a vector of the required quantiles. For example, if a plot of the 25th, 50th, and 75th quantiles are required, then `levels = c(0.25, 0.5, 0.75)`. Bivariate data have the option of being plotted as an intensity map instead of scatter plot. This can be obtained by setting `map="heat"`. There is also an option for plotting a cluster map of a fitted model using the option `map="cluster"`. Plots for specific components of a mixture model can be requested with the argument `component`. When `component=NULL` (which is default), the mixture contour is plotted. When `component` is a vector with length between 1 and g , the specified components are plotted and the mixing proportion is not taken into account. The last argument of the `fmmst.contour` functions “...” allows the user to pass additional arguments to the plot function, such as the colour and size of the points.

Figure 1a shows the contour of the fitted MST model to the Lymphoma data. Here, a heatmap of the original data is used. This plot can be generated via the command,

```

R> fmmst.contour.2d(Lympho, model=Fit, map="heat",
  xlab="SLP76", ylab="ZAP70")

```

The default `fmmst.contour.2d` function will return a scatter plot of the data in 2D superimposed with the contours of the fitted mixture model. For example, the following command generates a contour plot of the fitted FM-uMST model to the `ais` data in Section 4.2 (Figure 1b). Note that `fmmst.contour.2d` coloured the sample points according to the clustering given by the argument `clusters`:

```

R> label <- ais$Sex
R> label[label==0] <- 2
R> fmmst.contour.2d(ais[,c(2,12)], model=Fit2, clusters=label,
  xlab="Ht", ylab="Bfat")

```

Suppose we are interested in visualizing a clustering map of the fitted model to the simulated data in Section 4.4. This plot can be generated by issuing the following command.

```

R> fmmst.contour.2d(X, model=trainmodel, clusters=X[,3], map="cluster",
  component=1:3)

```




Figure 1: 2D contour plots generated by the `fmmst.contour.2d` function. (a) The fitted contour of the single component uMST model plotted over the hue intensity diagram of the Lymphoma dataset; (b) the default mixture contour plot of the fitted two-component FM-uMST model of the AIS dataset; (c) the contour of the individual components of the three-component model fitted to a bivariate synthetic sample plotted over the cluster map of the sample.

The output is given in Figure 1c. Note that the grid size of the cluster map is also controlled by the argument `grid` whose default value is 50.

To demonstrate the use of `fmmst.contour.3d`, we consider the clustering of a trivariate Diffuse Large B-cell Lymphoma (DLBCL) dataset provided by the British Columbia Cancer Agency. The data contain over 8000 cells derived from the lymph nodes of patients diagnosed with DLBCL. The samples were stained with three markers CD3, CD5, and CD19. The task is to cluster the cells into four groups. Hence we fit a four-component FM-uMST model to the data. The maximum number of iterations was increased to 300.

A scatterplot of the data is shown in Figure 2, where the dots are coloured according to the clustering provided by human experts, which is taken as the ‘true’ class labels. Figure 2b shows the 95th percentile density contours of the four components of the fitted model which are displayed with matching colours. The 3D plot uses the `rgl` visualization device system, and hence supports user friendly interactive navigation. The plots can be rotated in real-time to select a suitable viewpoint. The following code can be used to generate the 3D plots in Figure 2.

```
R> Fit4 <- fmmst(DLBCL, 4, itmax=300)
R> fmmst.contour.3d(DLBCL, model=Fit4, level=0.9, drawpoints=FALSE,
  xlab="FL1.LOG", ylab="FL2.LOG", zlab="FL4.LOG", quantile=0.95)
```

Results from manual expert gating enable us to calculate an error rate of clustering, which is measured by choosing among the possible permutations of cluster labels the one that gives the best performance. Note that dead cells were removed before evaluating the error rate against the benchmark results. For comparison, we calculated the error rate associated with the clustering results given by three other methods – FLAME (Pyne *et al.* 2009a), flowClust (Lo *et al.* 2008) and flowMeans (Aghaeepour *et al.* 2011). From Table 2, the FM-uMST model

Method	FLAME	flowClust	flowMeans	FM-uMST
Error rate	0.074	0.076	0.140	0.046

Table 2: Error rate of misclassification of four methods for the DLBCL dataset.




Figure 2: 3D contours plot of the DLBCL dataset generated by the `fmmst.contour.3d` function. (a) A scatterplot of the data coloured according to the the true clustering labels of the DLBCL dataset; (b) fitted contour of the three component FM-uMST model for the DLBCL dataset.

clearly shows superior performances in this dataset.

5. Concluding remarks

We have presented the R package **EMMIX-uskew** for fitting finite mixtures of unrestricted multivariate skew t -distributions to heterogeneous asymmetric data. The package implements a closed-form EM algorithm for fitting FM-uMST models and provides user-friendly visualization of the fitted contours in 2D and 3D. The major features of the software have been demonstrated on three real examples on the T-cell phosphorylation data, the Australian Institute of Sports (AIS) data and the DLBCL dataset. The clustering results were compared to mixture models with restricted multivariate skew t components and other methods. In both the AIS and DLBCL illustrations, the unrestricted model gave better clustering results with respect to the true class labels.

Acknowledgments

This work is supported by a grant from the Australian Research Council. Also, we would like to thank Professor Seung-Gu Kim for comments and corrections, and Drs. Kui (Sam) Wang and Saumyadipta Pyne for their helpful discussions on this topic.

References

- Aghaeepour N, Nikolov R, Hoos H, Brinkman R (2011). “Rapid cell population identification in flow cytometry data.” *Cytometry. Part A : the journal of the International Society for Analytical Cytology*, **79**, 6–13.
- Azzalini A (1985). “A class of distributions which includes the normal ones.” *Scandinavian Journal of Statistics*, **12**, 171–178.
- Azzalini A, Dalla Valle A (1996). “The multivariate skew-normal distribution.” *Biometrika*, **83**(4), 715–726.
- Banfield JD, Raftery A (1993). “Model-based Gaussian and non-Gaussian clustering.” *Biometrics*, **49**, 803–821.
- Basso RM, Lachos VH, Cabral CRB, Ghosh P (2010). “Robust mixture modeling based on scale mixtures of skew-normal distributions.” *Computational Statistics and Data Analysis*, **54**, 2926–2941.
- Böhning D (2000). *Computer Assisted Analysis of Mixtures and Applications: Meta-Analysis, Disease Mapping, and Others*. Chapman and Hall/CRC, London.
- Cabral C, Lachos V, Prates M (2012). “Multivariate mixture modeling using skew-normal independent distributions.” *Computational Statistics and Data Analysis*, **56**, 126–142.
- Cook RD, Weisberg S (1994). *An Introduction to Regression Graphics*. John Wiley & Sons, New York.
- Everitt BS, Hand DJ (1981). *Finite Mixture Distributions*. Chapman and Hall, London.
- Fraley C, Raftery AE (1999). “How many clusters? Which clustering methods? Answers via model-based cluster analysis.” *Computer Journal*, **41**, 578–588.
- Frühwirth-Schnatter S (2006). *Finite Mixture and Markov Switching Models*. Springer-Verlag, London.
- Frühwirth-Schnatter S, Pyne S (2010). “Bayesian inference for finite mixtures of univariate and multivariate skew-normal and skew- t distributions.” *Biostatistics*, **11**, 317–336.
- Gupta AK (2003). “Multivariate skew- t distribution.” *Statistics*, **37**, 359–363.
- Ho H, Lin T, Chen H, Wang W (2012). “Some results on the truncated multivariate t distribution.” *Journal of Statistical Planning and Inference*, **142**, 25–40.
- Lachos VH, Ghosh P, Arellano-Valle RB (2010). “Likelihood based inference for skew normal independent linear mixed models.” *Statistica Sinica*, **20**, 303–322.
- Lee S, McLachlan G (2011). “On the fitting of mixtures of multivariate skew t -distributions via the EM algorithm.” *arXiv:1109.4706 [stat.ME]*.
- Lee S, McLachlan G (2012a). “Finite mixtures of multivariate skew t -distributions: some recent and new results.” *Statistics and Computing*, **To appear**.

- Lee S, McLachlan G (2012b). “On mixtures of skew-normal and skew t -distributions.” *arXiv:1211.3602 [stat.ME]*.
- Lin TI (2010). “Robust mixture modeling using multivariate skew t distribution.” *Statistics and Computing*, **20**, 343–356.
- Lindsay BG (1995). *Mixture Models: Theory, Geometry, and Applications*. NSF-CBMS Regional Conference Series in probability and Statistics, Volume 5, Institute of Mathematical Statistics, Hayward, CA.
- Lo K, Brinkman R, Gottardo R (2008). “Automated gating of flow cytometry data via robust model-based clustering.” *Cytometry. Part A : the journal of the International Society for Analytical Cytology*, **73**, 321–332.
- Maier LM, Anderson DE, De Jager PL, Wicker L, Hafler DA (2007). “Allelic variant in CTLA4 alters T cell phosphorylation patterns.” *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 18607–18612.
- McLachlan GJ, Basford KE (1988). *Mixture Models: Inference and Applications*. Marcel Dekker, New York.
- McLachlan GJ, Peel D (2000). *Finite Mixture Models*. Wiley Series in Probability and Statistics.
- Mengersen K, Robert C, Titterton M (2011). *Mixtures: Estimation and Applications*. John Wiley & Sons, US.
- Prates M, Lachos V, Cabral C (2011). *mixsmsn: Fitting finite mixture of scale mixture of skew-normal distributions*. R package version 0.3-2, URL <http://CRAN.R-project.org/package=mixsmsn>.
- R Development Team (2011). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Pyne S, Hu X, Wang K, Rossin E, Lin TI, Maier LM, Baecher-Allan C, McLachlan GJ, Tamayo P, Hafler DA, De Jager PL, Mesirov JP (2009a). “Automated high-dimensional flow cytometric data analysis.” *PNAS*, **106**, 8519–8524.
- Pyne S, Hu X, Wang K, Rossin E, Lin TI, Maier LM, Baecher-Allan C, McLachlan GJ, Tamayo P, Hafler DA, De Jager PL, Mesirov JP (2009b). *FLAME: Flow analysis with Automated Multivariate Estimation*. URL http://www.broadinstitute.org/cancer/software/genepattern/modules/FLAME/published_data.
- Sahu S, Dey D, Branco M (2003). “A New Class of Multivariate Skew Distributions with Applications to Bayesian Regression Models.” *The Canadian Journal of Statistics*, **31**, 129–150.
- Titterton DM, Smith AFM, Markov UE (1985). *Statistical analysis of finite mixture distributions*. John Wiley & Sons, New York.

Wang K (2009). *EMMIX-skew: EM Algorithm for Mixture of Multivariate Skew Normal/ t Distributions*. R package version 1.0-12, URL http://www.maths.uq.edu.au/~gjm/mix_soft/EMMIX-skew.

Affiliation:

Geoffrey J. McLachlan
Department of mathematics
University of Queensland
Brisbane, Australia
E-mail: g.mclachlan@uq.edu.au